



333 South Kirkwood Road, Suite 200
Kirkwood, MO 63122

Office (314) 991-4335 Fax (314) 991-4340 Exchange: 888-456-8166

Nathan A. Mall, MD

Collin Magilligan, NP

Post-Operative Protocol- Multidirectional Stabilization

0-6 Weeks:

- Slingshot / Gunslinger Brace for 6 weeks
- Isometrics in brace, gentle supported Codman exercises
- PROM only for 6 weeks
- Grip strengthening, elbow ROM, wrist ROM

6-12 Weeks:

- Sling at night, can discontinue using the sling during the day
- AROM only as tolerated to increase ROM; no PT stretching or manipulation
- Restrict to 140° FF/ 40° ER at side/ IR to stomach/ 45° Abduction
- Scapular stabilization exercises avoiding anterior capsule stress
- Begin light isometrics for rotator cuff and deltoid, with arm at the side
- Can begin stationary bike

3-12 Months:

- Advance strengthening as tolerated: isometrics → bands → light weights (1-5 lbs); 8-12 reps/2-3 set per exercise for rotator cuff, deltoid, and scapular stabilizers
- Only do strengthening 3x/week to avoid rotator cuff tendonitis
- If ROM lacking, increase to full with gentle passive stretching at end ranges
- Begin eccentric motions, plyometrics (ex. Weighted ball toss), and closed chain exercises at 16 weeks
- Begin sports related rehab at 4 ½ months, including advanced conditioning
- Return to throwing at 6 months